

Rates of Comorbidities in Psoriatic Arthritis



Chronic inflammation in psoriatic arthritis can involve multiple systems and may contribute to the development of comorbidities.¹

48%^{2,3,*}

Obesity

24%¹

Hyperlipidemia

19%¹

Cardiovascular disease

34%¹

Hypertension

17%⁴

Depression

29%¹

Metabolic syndrome

13%¹

Diabetes

3%¹

Liver disease

*The National Health and Nutrition Examination Survey (NHANES) is a nationally representative survey of the US civilian, non-institutionalized population conducted by the CDC National Center for Health Statistics (NCHS). The cross-sectional survey includes an in-home interview to obtain sociodemographic characteristics and medical history, and a physical examination and laboratory measures, including BMI, taken at a mobile examination center. Patients self-reported being diagnosed with psoriatic arthritis.
1. Gupta S, et al. *Rheumatol Int*. 2021;41(2):275-284. 2. CDC NHANES Questionnaires, Datasets, and Related Documentation. <https://www.cdc.gov/nchs/nhanes/Default.aspx> (2009-2020). Accessed January 30, 2024. 3. Data on file. Lilly USA, LLC. DOF-IX-US-0341. 4. Zusman EZ, et al. *Semin Arthritis Rheum*. 2020;50(6):1481-1488.
VV-MED-162877 © 2024 Lilly USA, LLC. All rights reserved.